





دانشگاه زابل

مدیریت تحصیلات تکمیلی

دانشکده کشاورزی

گروه گیاهپزشکی

پایان نامه جهت اخذ درجه کارشناسی ارشد در رشته حشره شناسی کشاورزی

اثرات نانو فرمولاسیون ایمیداکلوپراید و تیمتوکسام بر بقا و بیان ژن های ضد میکروبی زنبورعسل

استاد راهنما :

دکتر نجمه صاحب زاده

استاد مشاور:

دکتر منصوره رخشانی پور

تهیه و تدوین :

مطهره راستگو

دی ۱۴۰۲

زنبورهای عسل یکی از گرده افشان‌های اصلی گیاهان هستند که در هنگام فعالیت جستجوگری در مزارع و باغات با سموم کشاورزی بکار برده شده مواجه می‌شوند. با افزایش تقاضای محصولات کشاورزی، استفاده از آفتکش‌های تجاری و نانوفرمولاسیون‌های آن‌ها برای از بین بردن آفات نیز افزایش یافته است. در بین آفتکش‌های مختلف، اثرات حاد و مزمن سموم نئونیکوتینوئیدی بر فعالیت زنبورعسل به خوبی مطالعه شده اما به بررسی اثرات نانوفرمولاسیون‌های این سموم به دلیل تنوع وسیع سنتز آنها بر بقا و فیزیولوژی با تاکید بر سیستم ایمنی زنبورعسل، پرداخته نشده است. با در نظر گرفتن این هدف، فرمولاسیون‌های نانوایمیداکلوپراید و نانوتیامتوکسام تهیه و زیست‌سنجی مقایسه‌ای آنها با فرمولاسیون‌های تجاری مربوطه هر کدام از آفتکش‌ها روی زنبورهای عسل جوان همسن بررسی شد. خصوصیات فیزیکی نانوفرمولاسیون‌های سنتزی با میکروسکوپ الکترونی روبشی (FESEM)، طیف‌سنجی مادون قرمز (FTIR) و پراش پرتو ایکس آنالیز (XRD) بررسی و بارگذاری نانوذرات با آفتکش‌های نامبرده مورد تأیید قرار گرفت. نتایج آنالیز FESEM نشان داد که نانوفرمولاسیون‌ها دارای کمترین تراکم و انباشتگی بودند. در آنالیز FTIR، پیک‌های شاخص تعیین شد. نتایج XRD نشان داد که اندازه تقریبی ذرات نانوایمیداکلوپراید و نانوتیامتوکسام به ترتیب ۶۱/۶ و ۵۶ نانومتر بود. در pH 5.5 میزان بارگذاری سموم در حامل (کیتوزان سدیم - آلژینات) افزایش معنی‌داری نشان داد. در این pH نیز درصد رهاسازی هر دو سم از حامل تا ۲۴ ساعت ادامه داشت. زیست‌سنجی با دو روش خوراکی و تماسی انجام شد. نتایج زیست‌سنجی‌ها نشان داد که میزان سمیت با سنتز فرم نانوی سموم مورد مطالعه کاهش داشت. همچنین میزان بقای زنبورهای عسل بررسی و مشخص شد که فرمولاسیون‌های تجاری اثر منفی بیشتری در بقای زنبورها نسبت به نانوفرمولاسیون‌ها داشتند. میزان بیان ژنهای *defensin2* و *abaecin* با کاهش غلظت سم در زنبورعسل کاهش یافت که می‌تواند بطور غیرمستقیم نشانگر مداخله در سیستم ایمنی زنبورعسل باشند.

واژه‌های کلیدی: بیان ژن، زنبورعسل ایرانی، زیست‌سنجی، مرگ و میر، نانوفرمولاسیون، نئونیکوتینوئیدها.

Abstract

Honey bees are among the primary pollinators of plants that encounter agricultural pesticides during foraging in fields and orchards. By the increasing demand for agricultural products, commercial pesticides, and nanoformulations severely apply to eradicate pests. Chronic and acute effects of pesticides, including neonicotinoids, have been well studied on honey bee activities. Still, due to their wide variety in synthesis, the assessment of nanoformulations of these pesticides on the survival and physiology, emphasizing the immune system of honey bees, has not been addressed. With this objective in mind, nanoformulations of imidacloprid and thiamethoxam were prepared. A comparative bioassay study of these formulations with their respective commercial formulations was conducted on young honey bees (1 to 3 days old). Before the bioassay assessment, the physical properties of synthetic nanoformulations were examined using Field Emission Scanning Electron Microscopy (FESEM), Fourier Transform Infrared Spectroscopy (FTIR), and X-ray Diffraction (XRD) analyses to confirm the loading of nanoparticles with the mentioned pesticides. The FESEM analysis revealed that the nanoformulations exhibited the lowest density and aggregation. In the FTIR analysis, characteristic peaks for nano-imidacloprid were observed at 3476.98, 3298.96, 29828.99, 1566.98, and within the 1300-1200 cm range. For nano-thiamethoxam, peaks appeared at 3286.96, 2886.96, 1586.64, and 1264.03 cm. XRD analysis results indicated approximate particle sizes for nano-imidacloprid and nano-thiamethoxam at 61.6 and 56 nanometers, respectively. At pH 5.5, the loading capacity of each pesticide (separately) into the studied carrier (sodium chitosan-alginate) showed a significant increase. At this pH, the release percentage of both pesticides from the carrier continued for up to 24 hours. Bioassay tests were conducted using two methods: oral and contact. The bioassay tests indicated reduced toxicity levels with nano-sized formulations of pesticides. Additionally, honeybee survival rates were examined in comparative treatments between commercial pesticide formulations and nano-formulations. It was determined that commercial formulations negatively impacted honeybee survival more than nano-formulations. Moreover, the expression levels of genes *abaecin* and *defensin2* upregulated with a reduction in pesticide concentration in honeybees. This indirectly suggests they could indicate interference within the honeybee's immune system.

Keywords: Cellular immunity, humoral immunity, antimicrobial peptides, Iranian honey bee, mortality



University of Zabol

Graduate School

Faculty of Plant Protection

The Thesis Submitted for The Degree of M.Sc

(In The field of Agricultural Entomology)

**Effects of imidacloprid and thiamethoxam nanoformulations on
the survival and expression of antimicrobial genes of
honeybees**

Supervisor:

Dr. N. Sahebzadeh

Advisor:

Dr. M. Rakhshanipour

By:

M. Rastgou

December 2023